

ESTIMATION AND APPLICATIONS OF QUANTILE REGRESSION FOR BINARY LONGITUDINAL DATA

Mohammad Arshad Rahman^a and Angela Vossmeier^b

^aIndian Institute of Technology Kanpur, India

^bClaremont McKenna College, USA

ABSTRACT

This chapter develops a framework for quantile regression in binary longitudinal data settings. A novel Markov chain Monte Carlo (MCMC) method is designed to fit the model and its computational efficiency is demonstrated in a simulation study. The proposed approach is flexible in that it can account for common and individual-specific parameters, as well as multivariate heterogeneity associated with several covariates. The methodology is applied to study female labor force participation and home ownership in the United States. The results offer new insights at the various quantiles, which are of interest to policymakers and researchers alike.

Keywords: Bayesian inference; binary outcomes; female labor force participation; home ownership; limited dependent variables; panel data

1. INTRODUCTION

The proliferation of panel data studies is well-documented and much of it has been attributed to data availability and challenging methodology (Hsiao, 2007). While panel data have been attractive for understanding behavior and dynamics, the modeling complexities involved have moved attention away from the data's

Topics in Identification, Limited Dependent Variables, Partial Observability,
Experimentation, and Flexible Modeling

Advances in Econometrics, Volume 40B, 157–191

Copyright © 2019 by Emerald Publishing Limited

All rights of reproduction in any form reserved

ISSN: 0731-9053/doi:10.1108/S0731-90532019000040B009

unique capacities. Modeling features such as a binary outcome variable or a quantile analysis, which are relatively straightforward to implement with cross-sectional data, are challenging and computationally burdensome for panel data. However, these features are important as they allow for the modeling of probabilities and lead to a richer view of how the covariates influence the outcome variable. Motivated by these difficulties, this chapter adds to the methodological advancements for panel data by developing quantile regression methods for binary longitudinal data and designing a computationally efficient estimation algorithm. The approach is applied to two empirical studies, female labor force participation and home ownership.

The chapter touches on three growing econometric literatures – discrete panel data, quantile regression for panel data, and quantile regression for discrete data. In reference to the latter, quantile regression has been implemented in binary data models (Benoit & Poel, 2012; Kordas, 2006), ordered data models (Alhamzawi & Ali, 2018; Rahman, 2016), count data models (Machado & Silva, 2005), and censored data models (Harding & Lamarche, 2012; Portnoy, 2003). For limited dependent variables, the concern is modeling the latent utility differential in the quantile framework, since the response variable takes limited values and does not yield continuous quantiles. Our chapter follows the work in this literature by using the latent utility setting and interpreting the utility as a “propensity” or “willingness” that underlie the latent scale, thus increasing our understanding of the impact of the covariates on the binary outcomes.

The literature on quantile regression in panel data settings includes (but is not limited to) Koenker (2004), Geraci and Bottai (2007), Y. Liu and Bottai (2009), Galvao (2010), Galvao and Kato (2016), Lamarche (2010), Harding and Lamarche (2009), and Harding and Lamarche (2017). The latter of these papers discusses the issues associated with solely focusing on fixed effects estimators and highlights the usefulness of allowing for a flexible specification of individual heterogeneity associated with covariates, also of interest in the present chapter. In a recent Bayesian paper, Luo, Lian, and Tian (2012) develop a hierarchical model to estimate the parameters of conditional quantile functions with random effects. The authors do so by adopting an asymmetric Laplace (AL) distribution for the residual errors and suitable prior distributions for the parameters. However, directly using the AL distribution does not yield tractable conditional densities for all of the parameters and hence a combination of Metropolis–Hastings (MH) and Gibbs sampling is required for model estimation. The use of the MH algorithm may require tuning at each quantile. To overcome this limitation, Luo et al. (2012) also present a full Gibbs sampling algorithm that utilizes the normal-exponential mixture representation of the AL distribution. This mixture representation is also followed in our work, with important computational improvements.

Finally, for discrete panel data, recent work by Bartolucci and Nigro (2010) introduces a quadratic exponential model for binary panel data and utilizes a conditional likelihood approach, which is computationally simpler than previous classical estimators. Bayesian approaches to binary panel data models include work by Albert and Chib (1996), Chib and Carlin (1999), Chib and Jeliazkov (2006),

and Burda and Harding (2013). These work influence the estimation methods designed in our quantile approach to binary panel data.

This chapter contributes to the three literatures by extending the various methodologies to a hierarchical Bayesian quantile regression model for binary longitudinal data (QBLD) and proposing a Markov chain Monte Carlo (MCMC) algorithm to estimate the model. The model handles both common (fixed) and individual-specific (random) parameters (commonly referred to as mixed effects in statistics). The algorithm implements a blocking procedure that is computationally efficient and the distributions involved allow for straightforward calculations of covariate effects. The framework is implemented in two empirical applications. The first application examines female labor force participation, which has been heavily studied in panel form. The topic became of particular interest in the state dependence versus heterogeneity debate (Heckman, 1981a). We revisit this question and implement our panel quantile approach, which has been otherwise unexplored for this topic. The results offer new insights regarding the determinants of female labor force participation and how the ages of children have different effects across the quantiles and utility scale. The findings suggest that policy should be focused on women's transitions into the labor force after child birth and the few years after.

The second application considers the probability of home ownership during the Great Recession. Microlevel empirical analyses on individuals moving into and out of housing markets are lacking in the recent literature. Past studies include Carliner (1974) and Poirier (1977), but the recent housing crisis offers a new opportunity to reevaluate the topic. Furthermore, a full quantile analysis of home ownership is yet to be explored. Since home ownership is a choice that requires years of planning, individual characteristics may range drastically across the latent utility scale. The analysis presented in this chapter controls for multivariate heterogeneity in individuals and wealth, and investigates the determinants of home ownership, state dependence in home ownership, and how the shock to housing markets affected these items. The results provide an understanding as to how individuals of particular demographics and socioeconomic status fared during the collapse of the housing market.

The rest of the chapter is organized as follows. Section 2 reviews quantile regression and the AL distribution, Section 3 introduces the QBLD model, presents a simulation study, and discusses methods for covariate effects. Section 4 considers the two applications and concluding remarks are offered in Section 5.

2. QUANTILE REGRESSION AND THE ASYMMETRIC LAPLACE DISTRIBUTION

The p -th quantile of a random variable Y is the value y_0 such that the probability that Y will be less than y_0 equals $p \in (0, 1)$. Mathematically, if $Q(\cdot)$ denotes

the inverse of the cumulative distribution function (*cdf*) of Y , the p -th quantile is defined as:

$$Q_Y(p) \equiv F_Y^{-1}(p) = \inf\{y_0 : F(y_0) \geq p\}.$$

Quantile regression implements the idea of quantiles within the regression framework with $Q(\cdot)$ modified to denote the inverse *cdf* of the dependent variable given the covariates. The objective is to estimate *conditional quantile functions* and to this purpose, regression quantiles are estimated by minimizing the quantile objective function which is a sum of asymmetrically weighted absolute residuals.

To formally explain the quantile regression problem, consider the following linear model,

$$y_i = x_i' \beta_p + \varepsilon_i, \quad \text{with} \quad Q_{\varepsilon_i}(p|x_i) = 0, \quad (1)$$

where y_i is a scalar response variable, x_i is a $k \times 1$ vector of covariates, β_p is a $k \times 1$ vector of unknown parameters that depend on quantile p , and ε_i is the error term such that its p -th quantile equals zero. Henceforth, we will drop the subscript p for notational simplicity. In classical econometrics, the error ε does not (or is not assumed to) follow any distribution and estimation requires minimizing the following objective function,

$$\min_{\beta \in \mathbf{R}^k} \left[\sum_{i: y_i < x_i' \beta} (1-p)|y_i - x_i' \beta| + \sum_{i: y_i \geq x_i' \beta} p|y_i - x_i' \beta| \right]. \quad (2)$$

The minimizer $\hat{\beta}$ gives the p -th regression quantile and the estimated conditional quantile function is obtained as $\hat{y}_i = x_i' \hat{\beta}$. Alternatively, the objective function in Eq. (2) can be written as a sum of piecewise linear or check functions as follows:

$$\min_{\beta \in \mathbf{R}^k} \sum_{i=1}^n \rho_p(y_i - x_i' \beta),$$

where $\rho_p(u) = u \cdot (p - I(u < 0))$ and $I(\cdot)$ is an indicator function, which equals one if the condition inside the parenthesis is true and zero otherwise. The check function, as seen in Fig. 1, is not differentiable at the origin. Hence, classical econometrics relies on computational techniques to estimate quantile regression models. Such computational methods include the simplex algorithm (Barrodale & Roberts, 1973; Dantzig, 1963; Dantzig & Thapa, 1997, 2003; Koenker & d'Orey, 1987), the interior point algorithm (Karmarkar, 1984; Mehrotra, 1992; Portnoy & Koenker, 1997), the smoothing algorithm (C. Chen, 2007; Madsen & Nielsen, 1993), and metaheuristic algorithms (Rahman, 2013).

In contrast to classical quantile regression, Bayesian quantile regression assumes that the error follows an AL distribution because the AL probability density function (*pdf*) contains the quantile loss function (Eq. 2) in its exponent.




Fig. 1. Quantile Regression Check Function.

This facilitates the construction of a working likelihood, required for Bayesian analysis. Maximizing an AL likelihood is equivalent to minimizing the quantile objective function (Koenker & Machado, 1999; Yu & Moyeed, 2001). A random variable Y follows an AL distribution if its *pdf* is given by:

$$f(y|\mu, \sigma, p) = \frac{p(1-p)}{\sigma} \exp\left[-\rho_p\left(\frac{y-\mu}{\sigma}\right)\right], \tag{3}$$

where $\rho_p(\cdot)$ is the check function as defined earlier, $-\infty < \mu < \infty$ is the location parameter, $\sigma > 0$ is the scale parameter, and $0 < p < 1$ is the skewness parameter (Kotz, Kozubowski, & Podgorski, 2001; Yu & Zhang, 2005). The mean and variance of Y from the *pdf* given by Eq. (3) are:

$$E(Y) = \mu + \frac{\sigma(1-2p)}{p(1-p)} \quad \text{and} \quad V(Y) = \frac{\sigma^2(1-2p+2p^2)}{p^2(1-p)^2}.$$

If $\mu = 0$ and $\sigma = 1$, then both mean and variance depend only on p and hence are fixed for a given value of p .

The Bayesian approach to quantile regression for binary data assumes that $\varepsilon \sim AL(0, 1, p)$. Here, the variance is constant to serve as a normalization for identification, typical in probit and logit models (Jeliakov & Rahman, 2012; Koop & Poirier, 1993; Poirier & Ruud, 1988). However, working directly with the AL distribution is not conducive to constructing a Gibbs sampler and hence the normal-exponential mixture of the AL distribution is often employed (Kozumi & Kobayashi, 2011). Several recent papers have utilized the mixture representation, including Ji, Lin, and Zhang (2012) for Bayesian model selection in binary and Tobit quantile regression, Luo et al. (2012) for estimating linear longitudinal data models, and Rahman (2016) for estimating ordinal quantile regression models. We also exploit the normal-exponential mixture representation of the AL distribution to derive the estimation algorithm for quantile regression in binary longitudinal data settings.

3. THE QUANTILE REGRESSION MODEL FOR BINARY LONGITUDINAL DATA

This section presents the QBLD model and an estimation algorithm to fit the model. The performance of the proposed algorithm is illustrated in a simulation study. The last part of this section considers methods for model comparison and covariate effects.

3.1. The Model

The proposed model looks at quantiles of binary longitudinal data expressed as a function of covariates with common effects and individual-specific effects. The individual-specific effects offer additional flexibility in that both intercept and slope heterogeneity can be captured, which are important to avoid biases in the parameter estimates. The QBLD model can be conveniently expressed in the latent variable formulation (Albert & Chib, 1993) as follows:

$$\begin{aligned} z_{it} &= x'_{it}\beta + s'_{it}\alpha_i + \varepsilon_{it}, \quad \forall i = 1, \dots, n, \quad t = 1, \dots, T_i, \\ y_{it} &= \begin{cases} 1 & \text{if } z_{it} > 0, \\ 0 & \text{otherwise,} \end{cases} \end{aligned} \quad (4)$$

where the latent variable z_{it} denotes the value of z at the t -th time period for the i -th individual, x'_{it} is a $1 \times k$ vector of explanatory variables, β is $k \times 1$ vector of common parameters, s'_{it} is a $1 \times l$ vector of covariates that have individual-specific effects, α_i is an $l \times 1$ vector of individual-specific parameters, and ε_{it} is the error term assumed to be independently and identically distributed (*i.i.d.*) as $AL(0, 1, p)$ with $Q_{\varepsilon_{it}}(p|x_{it}, \alpha_i) = 0$. This implies that the conditional density of $z_{it}|\alpha_i$ is an $AL(x'_{it}\beta + s'_{it}\alpha_i, 1, p)$ for $i = 1, \dots, n$, and $t = 1, \dots, T_i$, with $Q_{z_{it}}(p|x_{it}, \alpha_i) = x'_{it}\beta + s'_{it}\alpha_i$. Note that s_{it} may contain a constant for intercept heterogeneity, as well as other covariates (which are often a subset of those in x_{it}) to account for slope heterogeneity of those variables. The variable z_{it} is unobserved and represents the latent utility associated with the observed binary choice y_{it} . The latent variable formulation serves as a convenient tool in the estimation process. Furthermore, latent utility underlies the interpretation of the results at the various quantiles.

While working directly with the AL density is an option, the resulting posterior will not yield the full set of tractable conditional distributions necessary for a Gibbs sampler. Thus, we utilize the normal-exponential mixture representation of the AL distribution, presented in Kozumi and Kobayashi (2011), and express the error as follows:

$$\varepsilon_{it} = \theta w_{it} + \tau \sqrt{w_{it}} u_{it}, \quad \forall i = 1, \dots, n; \quad t = 1, \dots, T_i, \quad (5)$$

where $u_{it} \sim N(0, 1)$ is mutually independent of $w_{it} \sim \mathcal{E}(1)$ with \mathcal{E} representing an exponential distribution and the constants $\theta = \frac{1-2p}{p(1-p)}$ and $\tau = \sqrt{\frac{2}{p(1-p)}}$. The

mixture representation gives access to the appealing properties of the normal distribution.

Longitudinal data models often involve a moderately large amount of data, so it is important to take advantage of any opportunity to reduce the computational burden. One such trick is to stack the model for each individual i (Hendricks, Koenker, & Poirier, 1979). We define $z_i = (z_{i1}, \dots, z_{iT_i})'$, $X_i = (x'_{i1}, \dots, x'_{iT_i})'$, $S_i = (s'_{i1}, \dots, s'_{iT_i})'$, $w_i = (w_{i1}, \dots, w_{iT_i})'$, $D_{\tau\sqrt{w_i}} = \text{diag}(\tau\sqrt{w_{i1}}, \dots, \tau\sqrt{w_{iT_i}})$, and $u_i = (u_{i1}, \dots, u_{iT_i})'$. Building on Eqs. (4) and (5), the resulting hierarchical model can be written as,

$$\begin{aligned}
 z_i &= X_i\beta + S_i\alpha_i + \theta w_i + D_{\tau\sqrt{w_i}}u_i, \\
 y_{it} &= \begin{cases} 1 & \text{if } z_{it} > 0, \\ 0 & \text{otherwise,} \end{cases} \\
 \alpha_i|\varphi^2 &\sim N_i(0, \varphi^2 I), \quad w_{it} \sim \mathcal{E}(1), \quad u_{it} \sim N(0, 1), \\
 \beta &\sim N_k(\beta_0, B_0), \quad \varphi^2 \sim IG(c_1/2, d_1/2),
 \end{aligned} \tag{6}$$

where we assume that α_i are identically distributed as a normal distribution. The last row represents the prior distributions with N and IG denoting the normal and inverse-gamma distributions, respectively. Here, we note that the form of the prior distribution on β holds a penalty interpretation on the quantile loss function (Koenker, 2004). A normal prior on β implies a ℓ_2 penalty and has been used in Yuan and Yin (2010), and Luo et al. (2012). One may also employ a Laplace prior distribution on β that imposes ℓ_1 penalization, as used in Alhamzawi and Ali (2018). While Alhamzawi and Ali (2018) also work with quantile regression for discrete panel data (ordered, in particular), our work contributes by considering multivariate heterogeneity (not just intercept heterogeneity), and introducing computational improvements outlined below.

By Bayes' theorem, we express the "complete joint posterior" density as proportional to the product of likelihood function and the prior distributions as follows:

$$\begin{aligned}
 \pi(\beta, \alpha, w, z, \varphi^2|y) &\propto \left\{ \prod_{i=1}^n f(y_i|z_i, \beta, \alpha_i, w_i, \varphi^2) \pi(z_i|\beta, \alpha_i, w_i) \pi(w_i) \pi(\alpha_i|\varphi^2) \right\} \pi(\beta) \pi(\varphi^2), \\
 &\propto \left\{ \prod_{i=1}^n \left[\prod_{t=1}^{T_i} f(y_{it}|z_{it}) \right] \pi(z_i|\beta, \alpha_i, w_i) \pi(w_i) \pi(\alpha_i|\varphi^2) \right\} \pi(\beta) \pi(\varphi^2),
 \end{aligned} \tag{7}$$

where the first line uses independence between prior distributions and second line follows from the fact that given z_{it} , the observed y_{it} is independent of all parameters because the second line of Eq. (6) determines y_{it} given z_{it} with

probability one. Substituting the distribution of the variables associated with the likelihood and the prior distributions in Eq. (7) yields the following expression:

$$\begin{aligned} \pi(\beta, \alpha, w, z, \varphi^2 | y) &\propto \left\{ \prod_{i=1}^n \prod_{t=1}^{T_i} [I(z_{it} > 0)I(y_{it} = 1) + I(z_{it} \leq 0)I(y_{it} = 0)] \right\} \left\{ \prod_{i=1}^n |D_{T, \sqrt{w_i}}^2|^{-\frac{1}{2}} \right\} \\ &\times \exp \left[-\frac{1}{2} \sum_{i=1}^n \left\{ (z_i - X_i \beta - S_i \alpha_i - \theta w_i)' D_{T, \sqrt{w_i}}^{-2} (z_i - X_i \beta - S_i \alpha_i - \theta w_i) \right\} \right] \\ &\times \exp \left(-\sum_{i=1}^n \sum_{t=1}^{T_i} w_{it} \right) (2\pi\varphi^2)^{-\frac{nl}{2}} \exp \left[-\frac{1}{2\varphi^2} \sum_{i=1}^n \alpha_i' \alpha_i \right] \\ &\times (2\pi)^{-\frac{k}{2}} |B_0|^{-\frac{1}{2}} \exp \left[-\frac{1}{2} (\beta - \beta_0)' B_0^{-1} (\beta - \beta_0) \right] \times (\varphi^2)^{-\left(\frac{c_1}{2} + 1\right)} \exp \left[-\frac{d_1}{2\varphi^2} \right]. \end{aligned} \quad (8)$$

The joint posterior density in Eq. (8) does not have a tractable form, and thus simulation techniques are necessary for estimation. Bayesian methods are increasing in popularity (Poirier, 2006), and this chapter takes the approach for a couple of reasons. First, with discrete panel data, working with the likelihood function is complicated because it is analytically intractable. The inclusion of individual-specific effects makes matters worse. Second, while numerical simulation methods are available for discrete panel data, they are often slow and difficult to implement (Burda & Harding, 2013). The availability of a full set of conditional distributions (which are outlined below) makes Gibbs sampling an attractive option that will be simple to implement, both conceptually and computationally.

We can derive the conditional posteriors of the parameters and latent variables by a straightforward extension of the estimation technique for the linear mixed-effects model presented in Luo et al. (2012). This is presented as Algorithm 2 in Appendix 1, which shows the conditional posterior distributions for the parameters and latent variables necessary for a Gibbs sampler. While this Gibbs sampler is straightforward, there is potential for poor mixing properties due to correlation between (β, α_i) and (z_i, α_i) . The correlation often arises because the variables corresponding to the parameters in α_i are often a subset of those in x_{it} . Thus, by conditioning these items on one another, the mixing of the Markov chain will be slow.

To avoid this issue, we develop an alternative algorithm which jointly samples (β, z_i) in one block within the Gibbs sampler. This blocked approach significantly improves the mixing properties of the Markov chain. The success of these blocking techniques can be found in J. S. Liu (1994), Chib and Carlin (1999), and Chib and Jeliazkov (2006). The details of our blocked sampler are described in Algorithm 1.¹ In particular, β is sampled marginally of α_i from a multivariate normal distribution. The latent variable z_i is sampled marginally of α_i from a

truncated multivariate normal distribution denoted by $TMVN_{B_i}$, where B_i is the truncation region given by $B_i = (B_{i1} \times B_{i2} \times \dots \times B_{iT_i})$ such that B_{it} is the interval $(0, \infty)$ if $y_{it} = 1$ and the interval $(-\infty, 0]$ if $y_{it} = 0$. To draw from a truncated multivariate normal distribution, we utilize the method proposed in Geweke (1991). This involves drawing from a series of conditional posteriors which are univariate truncated normal distributions. Previous work using this approach include Chib and Greenberg (1998) and Chib and Carlin (1999). The random effects parameter α_i is sampled conditionally on β, z_i from another multivariate normal distribution. The variance parameter φ^2 is sampled from an IG distribution and finally the latent weight w is sampled element-wise from a generalized inverse Gaussian (GIG) distribution (Dagpunar, 1988, 1989; Devroye, 2014).

Algorithm 1. (Blocked Sampling)

- Sample (β, z_i) in one block. The objects (β, z_i) are sampled in the following two sub-steps.

- (1) Let $\Omega_i = (\varphi^2 S_i S_i' + D_{\tau, \sqrt{w_i}}^2)$. Sample β marginally of α from $\beta|z, w, \varphi^2 \sim N(\tilde{\beta}, \tilde{B})$, where,

$$\tilde{B}^{-1} = \left(\sum_{i=1}^n X_i' \Omega_i^{-1} X_i + B_0^{-1} \right) \text{ and } \tilde{\beta} = \tilde{B} \left(\sum_{i=1}^n X_i' \Omega_i^{-1} (z_i - \theta w_i) + B_0^{-1} \beta_0 \right).$$

- (2) Sample the vector $z_i|y_i, \beta, w_i, \varphi^2 \sim TMVN_{B_i}(X_i \beta + \theta w_i, \Omega_i)$ for all $i = 1, \dots, n$, where $B_i = (B_{i1} \times B_{i2} \times \dots \times B_{iT_i})$ and B_{it} are the interval $(0, \infty)$ if $y_{it} = 1$ and the interval $(-\infty, 0]$ if $y_{it} = 0$. This is done by sampling z_i at the j -th pass of the MCMC iteration using a series of conditional posterior distribution as follows:

$$z_{it}^j | z_{i1}^j, \dots, z_{i(t-1)}^j, z_{i(t+1)}^j, \dots, z_{iT_i}^j \sim TN_{B_{it}}(\mu_{it-t}, \Sigma_{it-t}), \text{ for } t = 1, \dots, T_i,$$

where TN denotes a truncated normal distribution. The terms μ_{it-t} and Σ_{it-t} are the conditional mean and variance, respectively, and are defined as,

$$\begin{aligned} \mu_{it-t} &= x'_{it} \beta + \theta w_{it} + \Sigma_{t,-t} \Sigma_{-t,-t}^{-1} (z^j_{i,-t} + (X_i \beta + \theta w_i)_{-t}), \\ \Sigma_{it-t} &= \Sigma_{t,t} - \Sigma_{t,-t} \Sigma_{-t,-t}^{-1} \Sigma_{-t,t}, \end{aligned}$$

where $z^j_{i,-t} = (z^j_{i1}, \dots, z^j_{i(t-1)}, z^j_{i(t+1)}, \dots, z^j_{iT_i})$, $(X_i \beta + \theta w_i)_{-t}$ is a column vector with t -th element removed, $\Sigma_{t,t}$ denotes the (t, t) -th element of Ω_i , $\Sigma_{t,-t}$ denotes the t -th row of Ω_i with element in the t -th column removed, and $\Sigma_{-t,-t}$ is the Ω_i matrix with t -th row and t -th column removed.

- Sample $\alpha_i | z, \beta, w, \varphi^2 \sim N(\tilde{a}, \tilde{A})$ for $i = 1, \dots, n$, where,

$$\tilde{A}^{-1} = \left(S'_i D_{\tau\sqrt{w_i}}^{-2} S_i + \frac{1}{\varphi^2} I_l \right) \quad \text{and} \quad \tilde{a} = \tilde{A} \left(S'_i D_{\tau\sqrt{w_i}}^{-2} (z_i - X_i \beta - \theta w_i) \right).$$

- Sample $w_{it} | z_{it}, \beta, \alpha_i \sim GIG(0.5, \tilde{\lambda}_{it}, \tilde{\eta})$ for $i = 1, \dots, n$ and $t = 1, \dots, T_i$, where,

$$\tilde{\lambda}_{it} = \left(\frac{z_{it} - x'_{it}\beta - s'_{it}\alpha_i}{\tau} \right)^2 \quad \text{and} \quad \tilde{\eta} = \left(\frac{\theta^2}{\tau^2} + 2 \right).$$

- Sample $\varphi^2 | \alpha \sim IG(\tilde{c}_1/2, \tilde{d}_1/2)$, where $\tilde{c}_1 = (nl + c_1)$ and $\tilde{d}_1 = \left(\sum_{i=1}^n \alpha'_i \alpha_i + d_1 \right)$.

We end this section with a cautionary note on sampling from a truncated multivariate normal distribution, with the hope that it will be useful to researchers on quantile regression. In our algorithm above, we sample z_i from $TMVN_{B_i}(X_i\beta + \theta w_i, \Omega_i)$ using a series of conditional posteriors which are univariate truncated normal distributions. This method is distinctly different and should not be confused with sampling from a recursively characterized truncation region typically related to the Geweke–Hajivassiliou–Keane (GHK) estimator (Börsch-Supan & Hajivassiliou, 1993; Geweke, 1991; Hajivassiliou & McFadden, 1998; Keane, 1994).² The difference between the two samplers have been exhibited in Breslaw (1994) and carefully discussed in Jeliazkov and Lee (2010).

3.2. Simulation Study

This subsection evaluates the performance of the algorithm in a simulation study, where the data are generated from a model that has common effects and individual-specific effects in both the intercept and slopes. We estimate the quantile regression model for binary longitudinal data (QBLD) using our proposed blocked sampler (Algorithm 1) and the non-blocked sampler (Algorithm 2).

The data are simulated from the model $z_{it} = x'_{it}\beta + s'_{it}\alpha_i + \varepsilon_{it}$ where $t = 1, \dots, 10$ and $i = 1, \dots, 500$. For the parameters and covariates: $\beta = (-5, 6, 4)'$, $\alpha_i \sim N(0_2, I_2)$, $x'_{it} = (1, x_{2it}, x_{3it})$ with $x_{2it} \sim U(0, 1)$ and $x_{3it} \sim U(0, 1)$, $s'_{it} = (1, s_{2it})$ with $s_{2it} \sim U(0, 1)$. The error is generated from a standard AL distribution, $\varepsilon_{it} \sim AL(0, 1, p)$ for $p = 0.25, 0.5, 0.75$. Here, the notation $U(0, 1)$ denotes a standard uniform distribution. The binary response variable y_{it} is constructed by assigning one to all positive values of z_{it} and zero to all negative values of z_{it} . Since the values generated from an AL distribution are different at each quantile, the number of zeros and ones are also different at each quantile. In the simulation, the number of observations corresponding to zeros and ones for the 25th, 50th, and 75th quantiles are (1566, 3444), (2588, 2412), and (3536, 1464), respectively.

Table 1. Results from the QBLD Model in the Simulation Study.

	Blocked Sampling								
	25th Quantile			50th Quantile			75th Quantile		
	Mean	Std	IF	Mean	Std	IF	Mean	Std	IF
β_1	-5.33	0.22	4.55	-5.06	0.18	4.09	-5.08	0.24	4.10
β_2	6.16	0.28	4.38	5.96	0.22	3.87	6.16	0.27	4.11
β_3	4.34	0.24	3.86	3.88	0.19	3.66	3.88	0.23	3.21
φ^2	0.95	0.16	4.68	0.66	0.11	4.60	0.81	0.15	4.93
Non-blocked Sampling									
β_1	-5.32	0.22	5.94	-5.05	0.20	6.90	-5.07	0.23	6.63
β_2	6.15	0.27	6.05	5.95	0.23	6.57	6.15	0.26	6.69
β_3	4.35	0.24	5.52	3.88	0.20	5.40	3.88	0.23	5.34
φ^2	0.95	0.16	5.58	0.66	0.11	5.26	0.81	0.14	6.15

Note: The first panel presents the results from Algorithm 1 and the second panel presents the results from Algorithm 2. Mean is posterior mean, Std is posterior standard deviation and IF is inefficiency factor.


The posterior estimates of the model parameters are based on the generated data and the following independent prior distributions: $\beta \sim N(0_k, 10I_k)$ and $\varphi^2 \sim IG(10/2, 9/2)$. Table 1 reports the posterior means, standard deviations, and inefficiency factors calculated from 12,000 MCMC iterations after a burn-in of 3,000 iterations. The inefficiency factors are calculated using the batch-means method discussed in Greenberg (2012). The simulation exercise was repeated for various covariates, sample sizes, and common and individual-specific parameters, and the results do not change from this baseline case; hence they are not presented.

The posterior means of the regression coefficients in both of the samplers (blocked and non-blocked methods) are near the true values, $\beta = (-5, 6, 4)'$. Additionally, the standard deviations are small. Across each quantile, the number of zeros and ones varies, and the samplers perform well in each case. Furthermore, starting the algorithm at different values appears inconsequential, which is a benefit of the full Gibbs sampler.

Turning attention to the differences between the two algorithms, it is clear that the inefficiency factors from the blocked algorithm are much lower, suggesting better sampling performance and a nice mixing of the Markov chain. The advantages of the blocking procedure are more apparent from the autocorrelation in the MCMC draws at different lags. Table 2 presents the autocorrelation in MCMC draws at lag 1, lag 5, and lag 10. Looking at lag 10, the autocorrelation for the β s is in the range of 0.25 – 0.43 in the blocked algorithm, which is nearly half of 0.55 – 0.73, obtained from the non-blocked sampler. Recall that in our data generation process, we did not make the covariates in s_{it} a subset of those in x_{it} . Whereas in real-data exercises, it is typical for s_{it} to be a subset.

Table 2. Autocorrelation in MCMC Draws at Lag 1, Lag 5, and Lag 10.

	Blocked Sampling								
	25th Quantile			50th Quantile			75th Quantile		
	Lag 1	Lag 5	Lag 10	Lag 1	Lag 5	Lag 10	Lag 1	Lag 5	Lag 10
β_1	0.86	0.59	0.41	0.85	0.54	0.35	0.88	0.61	0.41
β_2	0.89	0.61	0.43	0.87	0.53	0.34	0.89	0.60	0.39
β_3	0.86	0.50	0.31	0.83	0.44	0.25	0.84	0.45	0.23
φ^2	0.93	0.73	0.54	0.92	0.70	0.51	0.93	0.75	0.58
	Non-blocked Sampling								
β_1	0.96	0.84	0.71	0.97	0.85	0.73	0.97	0.85	0.71
β_2	0.96	0.81	0.68	0.96	0.82	0.68	0.96	0.80	0.65
β_3	0.95	0.77	0.61	0.95	0.77	0.60	0.94	0.75	0.55
φ^2	0.92	0.76	0.63	0.92	0.74	0.59	0.93	0.79	0.68

**Fig. 2.** Trace Plots of the MCMC Draws at the 25th Quantile from Algorithm 1.

Therefore, we expect the benefits of the blocked sampler to be even more pronounced in real-data settings.

Finally, Fig. 2 presents the trace plots of the parameters at the 25th quantile for the blocked algorithm, which graphically demonstrate the appealing sampling. Given the computational efficiency with the blocking procedure, it is our preferred way for estimating QBLD models and will be used in the subsequent real data applications.

3.3. Additional Considerations

In this section, we briefly discuss methods for model comparison and computation of covariate effects. For model comparison, we follow standard techniques for longitudinal data models. Specifically, in the application sections, we provide the log-likelihood, conditional Akaike Information Criterion (AIC), and conditional Bayesian Information Criterion (BIC). The calculations for the conditional AIC and conditional BIC can be found in Greven and Kneib (2010) and Delattre, Lavielle, and Poursat (2014), respectively. This is a bit unusual for a Bayesian analysis; however, we want the results in our empirical applications to align with the classical work on the topics, such as Bartolucci and Farcomeni (2012). Thus, we follow the approaches so as to allow for better comparisons and cross-references.

For covariate effects, in general terms, we are interested in the average difference in the implied probabilities between the case when x_{1it} is set to the value x_{1it}^\dagger and x_{1it}^\ddagger . Given the values of the other covariates denoted x_{-1it}, s_{it} and those of the model parameters θ , one can obtain the probabilities $\Pr(y_{it} = 1|x_{1it}^\dagger, x_{-1it}, s_{it}, \theta)$ and $\Pr(y_{it} = 1|x_{1it}^\ddagger, x_{-1it}, s_{it}, \theta)$. Following from Jeliazkov, Graves, and Kutzbach (2008) and Jeliazkov and Vossmeier (2018), if one is interested in the distribution of the difference $\{\Pr(y_{it} = 1|x_{1it}^\dagger) - \Pr(y_{it} = 1|x_{1it}^\ddagger)\}$ marginalized over $\{x_{-1it}, s_{it}\}$ and θ given the data y , a practical procedure is to marginalize out the covariates using their empirical distribution, while the parameters are integrated out with respect to their posterior distribution. Formally, the goal is to obtain a sample of draws from the distribution,

$$\begin{aligned} & \{\Pr(y_{it} = 1|x_{1it}^\dagger) - \Pr(y_{it} = 1|x_{1it}^\ddagger)\} \\ &= \int \{\Pr(y_{it} = 1|x_{1it}^\dagger, x_{-1it}, s_{it}, \theta) - \Pr(y_{it} = 1|x_{1it}^\ddagger, x_{-1it}, s_{it}, \theta)\} \\ & \quad \times \pi(x_{-1it}, s_{it}) \pi(\theta|y) d(x_{-1it}, s_{it}) d\theta. \end{aligned}$$

The probabilities are computed from the AL *cdf*, allowing for straightforward interpretations of the covariate effects. Also, the procedure handles uncertainty stemming from the sample and estimation strategy. This approach is demonstrated in each of the following applications.

4. APPLICATIONS

4.1. Female Labor Force Participation

Modeling female labor force participation has been an important area of work in the economics and econometric literature for decades. The list of work is vast, but a partial list includes Heckman and MaCurdy (1980, 1982), Mroz (1987), Hyslop (1999), Arellano and Carrasco (2003), Chib and Jeliazkov (2006), Kordas (2006), Carro (2007), Bartolucci and Nigro (2010), and Eckstein and Lifshitz (2011).

Within the literature, several pertinent questions have been analyzed including the relationship between participation and age, education, fertility, and

permanent and transitory incomes. However, serial persistence in the decision to participate and its two competing theories – heterogeneity and state dependence – have been of substantive interest. Heterogeneity implies that females may differ in terms of certain unmeasured variables that affect their probability of labor force participation. If heterogeneity is not properly controlled, then past decisions may appear significant to current decisions leading to what is called spurious state dependence. In contrast, pure state dependence implies that dynamic effects of past participation genuinely affect current employment decisions. Consideration of heterogeneity and state dependence is important in modeling female labor force participation and can have economic implications as discussed in Heckman (1981a, 1981b), and Hsiao (2014, pp. 261–270). We reexamine the above-mentioned aspects using our proposed QBLD model. To our knowledge, this is the first attempt to analyze female labor force participation within a longitudinal quantile framework. So, what can we learn from a panel quantile approach? Of particular interest are the impacts of infants and children across the various quantiles. Understanding the differential effects across the latent utility scale can help shape female labor force policies, such as maternity leave and child care.

Before proceeding forward, we draw attention to Kordas (2006) who evaluated female labor force participation using cross-sectional data and smoothed binary regression quantiles. His results offer interesting insights across the quantiles, which further motivate our application and extension to transitions into and out of the labor force in the panel setting. We also follow his interpretation where the latent utility differential between working and not working may be interpreted as a “propensity” or “willingness-to-participate” (WTP) index.

The data for this study are taken from Bartolucci and Farcomeni (2012), which were originally extracted from the Panel Study of Income Dynamics (PSID) conducted by the University of Michigan. The data consist of a sample of $n = 1446$ females who were followed for the period 1987 to 1993 with respect to their employment status and a host of demographic and socioeconomic variables. The dependent variable in the model is *employment* status (= 1 if the individual is employed, = 0 otherwise) and the covariates include *age* (in 1986), *education* (number of years of schooling), *child 1–2* (number of children aged one to two, referred to the previous year), *child 3–5*, *child 6–13*, *child 14–*, *Black* (indicator for Black race), *income* of the husband (in US dollars, referred to the previous year), and *fertility* (indicator variable for birth of a child in a certain year). Lagged employment status is also included as a covariate to examine state dependence.

Table 3 presents summary statistics for the variables. The presentation of the table follows from Hyslop (1999), where statistics are broken up into subgroups of women that have worked zero years, seven years, or transitioned during the period. As one can see from the table, the average age in the sample is roughly 30; about 40% of the sample are employed throughout the entire period, 10% are not in the labor force throughout the entire period, 20% transition into or out of the labor force once, and 30% transition multiple times. Looking closely

Table 3. Sample Characteristics of the Female Labor Force Participation Data.

	Full Sample	Employed Seven Years	Employed Zero Years	Single Transition from Work	Single Transition to Work	Multiple Transitions
	(1)	(2)	(3)	(4)	(5)	(6)
Age	29.55 (4.61)	30.44 (4.34)	29.18 (4.51)	29.21 (4.77)	29.23 (4.62)	28.68 (4.73)
Education	13.14 (2.06)	13.33 (1.98)	12.68 (2.15)	13.20 (2.13)	13.01 (2.19)	13.08 (2.05)
Child 1–2	0.31 (0.53)	0.22 (0.45)	0.46 (0.60)	0.31 (0.53)	0.34 (0.57)	0.38 (0.57)
Child 3–5	0.37 (0.57)	0.27 (0.49)	0.56 (0.65)	0.32 (0.54)	0.50 (0.65)	0.42 (0.60)
Child 6–13	0.75 (0.92)	0.71 (0.87)	0.92 (1.00)	0.55 (0.81)	0.99 (1.03)	0.74 (0.94)
Child 14–	0.32 (0.67)	0.39 (0.72)	0.31 (0.71)	0.29 (0.69)	0.26 (0.61)	0.26 (0.60)
Black	0.24 (0.43)	0.27 (0.44)	0.26 (0.44)	0.19 (0.39)	0.21 (0.40)	0.22 (0.41)
Income/10,000	3.04 (2.60)	2.82 (1.82)	3.81 (5.28)	3.43 (3.14)	2.99 (2.04)	2.96 (1.89)
Fertility	0.07 (0.25)	0.04 (0.21)	0.08 (0.28)	0.10 (0.29)	0.05 (0.22)	0.09 (0.28)
Years worked						
0	10.30	–	100	–	–	–
1	5.33	–	–	20.00	9.03	8.25
2	6.29	–	–	7.86	12.26	14.39
3	6.36	–	–	12.14	10.97	13.68
4	8.64	–	–	11.43	17.42	19.34
5	9.34	–	–	13.57	23.23	18.87
6	13.76	–	–	35.00	27.10	25.47
7	39.97	100	–	–	–	–
Observations	1446	578	149	140	155	424

Note: The first panel presents the mean/proportion and standard deviations (in parenthesis) of the variables in the full and subsamples. The second panel displays the column percentages for the number of years worked and the third panel (i.e., last row) presents the number of observations in the full and the subsamples.

at the different variables for children, there is a decent amount of variation across the subgroups. For women who are employed zero years, the average values for *child 1–2* and *child 3–5* are 0.46 and 0.56, respectively. These numbers are more than double compared to that of women who are employed for all

the seven years. Further, as children age (*child 6–13*) more, women have a single transition to work. While these differences demonstrate some observed heterogeneity, unobserved heterogeneity still plays a role, which motivates further analysis. Particularly, a quantile setting will reveal information not available in the raw observed data by utilizing the latent scale as the WTP index.

The data are modeled following Eqs. (4) and (5) and the model (QBLD) is specified with a random intercept (i.e., s_{it} only includes a constant). We also estimate the probit model for binary longitudinal data (PBLD) using the algorithm presented in Koop, Poirier, and Tobias (2007) and Greenberg (2012) and identical priors for relevant parameters. The results for the QBLD and PBLD models are presented in Table 4 and are based on data for the years 1988–1993, since using a lagged dependent variable drops information for the year 1987. The reported posterior estimates are based on 12,000 MCMC draws after a burn-in of 3,000 draws and the following priors on the parameters: $\beta \sim N(0_k, 10I_k)$ and $\varphi^2 \sim IG(10/2, 9/2)$. Table 4 presents the posterior means, standard deviations, and inefficiency factors at the 25th, 50th, and 75th quantiles, and for the binary probit model. Furthermore, the log-likelihood, conditional AIC (Greven &

Table 4. Results from the Female Labor Force Participation Study.

	QBLD									PBLD		
	25th Quantile			50th Quantile			75th Quantile			Mean	Std	IF
	Mean	Std	IF	Mean	Std	IF	Mean	Std	IF			
Intercept	-3.11	0.21	4.59	-0.31	0.18	4.45	1.35	0.23	4.79	-0.08	0.07	2.79
Age [†]	0.03	0.01	2.27	0.01	0.01	2.38	-0.01	0.02	2.72	0.01	0.01	1.57
(Age [†]) ² /100	-0.23	0.26	1.96	-0.19	0.25	2.06	-0.13	0.33	2.59	-0.08	0.10	1.45
Education [†]	0.17	0.03	2.29	0.21	0.03	2.57	0.28	0.05	3.18	0.08	0.01	1.59
Child 1–2	-0.22	0.11	2.68	-0.28	0.11	2.84	-0.38	0.13	2.97	-0.12	0.04	1.67
Child 3–5	-0.55	0.10	2.89	-0.52	0.10	3.22	-0.56	0.12	2.91	-0.21	0.04	1.74
Child 6–13	-0.17	0.07	2.39	-0.18	0.07	2.59	-0.18	0.08	2.95	-0.07	0.02	1.58
Child 14–	-0.05	0.10	2.66	-0.02	0.10	2.89	-0.01	0.13	3.22	-0.01	0.04	1.71
Black	0.20	0.15	2.02	0.24	0.15	2.24	0.26	0.19	2.69	0.09	0.06	1.53
Income [†] /10,000	-0.13	0.03	3.03	-0.14	0.02	3.00	-0.18	0.03	3.52	-0.05	0.01	1.95
Fertility	-1.91	0.20	2.71	-2.06	0.20	2.90	-2.60	0.33	3.85	-0.72	0.07	1.67
Lag Employment	4.89	0.16	3.75	3.88	0.13	4.47	6.71	0.20	5.24	1.49	0.05	3.34
φ^2	1.42	0.35	6.36	1.39	0.33	6.16	2.12	0.50	7.12	0.33	0.05	4.97
Log-likelihood	-3115.72			-3127.38			-3146.68			-2887.91		
AIC	6257.45			6280.77			6319.36			5801.82		
BIC	6354.82			6378.14			6416.74			5899.20		

Note: [†]denotes variable minus the sample average. Mean is posterior mean, Std is posterior standard deviation, and IF is inefficiency factor.

Kneib, 2010) and conditional BIC (Delattre et al., 2014) are available for each model.

First, note that across the quantiles the inefficiency factors are low, implying a nice mixing of the Markov chain. These results, which were demonstrated in the simulation study, hold in empirical applications as well. Next, if we consider each quantile as corresponding to a different likelihood, then the 25th quantile has the lowest conditional AIC and conditional BIC. This result is not surprising since the unconditional probability of participation is around 70% in the sample. Our result also finds support in Kordas (2006), where he reports that the 30th conditional quantile would be the one most efficiently estimable.

The results for the education variable are positive, statistically different from zero, and show various incremental differences across the quantiles. Education is found to have stronger effects in the upper part of the latent index, which is expected since these are women who have a high utility for working and thus have obtained the requisite education. Regarding the state dependence versus heterogeneity debate, we find that employment is serially positively correlated, which is a consequence of state dependence. The effect gets incrementally larger as one moves up the latent utility scale. While we are controlling for individual heterogeneity with the random intercept, we still find evidence of state dependence. This result agrees with Bartolucci and Farcomeni (2012), who investigate the question with a latent class model. Other papers that find empirical evidence of strong state dependence effects include Heckman (1981a), Hyslop (1999), and Chib and Jeliazkov (2006).

To further understand the results, covariate effects are computed for several variables for the three quantiles and the PBLD model. The covariate effect calculations follow from Section 3.3 and the results are displayed in Table 5. Note that the 50th quantile results are similar to that of the PBLD model, which is to be expected. The covariate effect for education is calculated on the restricted sample of individuals with a high school degree (12 years of schooling). The effect that is computed is four additional years of schooling, implying a college degree. The effect for income is a discrete change by \$10,000, the effect for children is increasing the count by one, and for fertility it is a discrete change to the indicator variable.

Table 5. Covariate Effects in the Female Labor Force Participation Study.

	QBLD			PBLD
	25th	50th	75th	
Education	0.0523	0.0711	0.0633	0.0698
Child 1–2	–0.0160	–0.0212	–0.0206	–0.0254
Child 3–5	–0.0415	–0.0397	–0.0302	–0.0430
Child 6–13	–0.0123	–0.0133	–0.0098	–0.0146
Income	–0.0095	–0.0102	–0.0097	–0.0105
Fertility	–0.1672	–0.1747	–0.1335	–0.1627

The results show that the birth of a child in that year (*fertility*) reduces the probability that a woman works by 16.7 percentage points at the 25th quantile, 17.4 percentage points at the 50th quantile, and 13.3 percentage points at the 75th quantile. For individuals in the lower part of the latent index, having children aged one to two impacts their employment decision less than those at the upper quantiles. Perhaps, women with a low utility for working are less impacted by infants and toddlers because it is often a desire to stay home with the child for a few years. Whereas, women with a high utility for working face negative impacts because of the desire to enter the workforce.

The most pronounced negative effect of children occurs when the child is aged three to five. Often women temporarily exit the workforce until children are ready for pre-school and this result provides evidence of the difficulty women face reentering the workforce after several years of leave of absence (Drange & Rege, 2013). The finding is interesting from a policy standpoint. If policy is focused on increasing participation, offering more support in the years when the child is likely not breastfeeding but before kindergarten would be beneficial.

The covariate effect of a college degree is 5.2 to 7.1 percentage points across the quantiles, while husband's income is approximately -1 percentage point across the quantiles. Thus, a college degree increases the probability a woman works by about 6 percentage points, whereas an increase in family income only decreases the probability by 1 percentage point for every \$10,000. While many of these results align with existing findings, the behavior in the high and low quantiles presents useful information, which was otherwise unexplored in panel data.

4.2. Home Ownership

The recent financial crisis had major implications for home ownership in the United States. Fig. 3 displays the home ownership rates for the United States from the 1960s to 2017. These data were taken from the FRED website provided by the Federal Reserve Bank of St. Louis. The rate of home ownership rose in the late 1990s and early 2000s, but started to decline after 2007. The determinants of home ownership was reviewed in the 1970s (Carliner, 1974;




Fig. 3. Home Ownership Rates in the United States. *Source:* Data taken from FRED, provided by the Federal Reserve Bank of St. Louis.

Poirier, 1977). However, the recent crisis offers a unique event and shock to housing markets to reevaluate this topic.

The literature on home ownership has examined racial gaps (Charles & Hurst, 2002; Turner & Smith, 2009), wealth accumulation and income (Turner & Luea, 2009), mobility and the labor market (Fairlie, 2013; Ferreira, Gyourko, & Tracy, 2010), and tax policy (Hilber & Turner, 2014). However, unlike the labor force context, state dependence has only been lightly examined with regard to housing tenure.³ Given the large down payments and extensive mortgage processes typical in home ownership, state dependence is likely to be a key factor, as well as individual heterogeneity.

Furthermore, quantile analyses in the home ownership literature are lacking. The quantiles represent degrees of *willingness* or *utility* of owning a home. Owning a home in the United States usually requires an individual to produce a large upfront investment, a promising credit history, and a willingness to engage in 30-year mortgages, resulting in less liquidity. Given these requirements, interest lies in how the determinants of home ownership vary across the latent utility scale. Therefore, this chapter adds to the literature on the probability of home ownership by employing the QBLD model. The approach has several advantages, namely that we can control for multivariate heterogeneity, visit the state dependence versus heterogeneity argument in the housing context, and analyze willingness of home ownership across the quantiles.

The dataset is constructed from the PSID and consists of a balanced panel of 4,092 individuals observed for the years 2001, 2003, 2005, 2007, 2009, 2011, and 2013. The sample is restricted to individuals aged 25–65 who answered the relevant questions for the seven years and captures the period before, during, and after the Great Recession. The dependent variable is defined as follows:

$$y_{it} = \begin{cases} 1 & \text{home owner} \\ 0 & \text{not a home owner,} \end{cases} \quad (9)$$

for $i = 1, \dots, 4092$ and $t = 2003, 2005, 2007, 2009, 2011, 2013$ (2001 is dropped because it is a dynamic model). The covariates include demographics, marital status, employment, job industry, health insurance, education, socioeconomic status, lagged home ownership, and an indicator for after the recession (2009–2013). The model includes a random intercept and a random slope on an income-to-needs variable, which allows for individual heterogeneity and heterogeneity in income. Heterogeneity in income is an important control because a marginal increase in income could have a wide range of effects on the probability of owning a home, where for some the effect of income could be zero (perhaps, those who own their home freehold, or those who have no desire for ownership). Whereas, for others, increases in income could go directly into home ownership utility. Table 6 presents summary statistics for the variables. Once again, the presentation of the table follows from Hyslop (1999), where statistics are broken up into subgroups of people that have always been home owners, never been home owners, or transitioned during the period of interest.

Table 6. Sample Characteristics of the Home Ownership Data.

	Full Sample (1)	Owned Six Years (2)	Owned Zero Years (3)	Single Transition from Ownership (4)	Single Transition to Ownership (5)	Multiple Transitions (6)
Age of head (2003)	45.74 (13.51)	48.78 (12.56)	42.31 (13.82)	46.71 (15.48)	38.91 (11.63)	39.62 (12.29)
No. of children	0.78 (1.15)	0.71 (1.08)	0.82 (1.25)	0.83 (1.21)	0.85 (1.12)	1.00 (1.27)
Income/10,000	7.71 (10.53)	9.66 (11.18)	3.17 (3.33)	6.54 (16.28)	6.74 (5.71)	6.82 (10.15)
Inc–Needs Ratio	4.89 (7.38)	6.07 (7.51)	2.17 (2.30)	4.25 (14.01)	4.37 (3.74)	4.23 (5.66)
Net wealth/10,000	23.05 (128.08)	35.76 (167.12)	1.98 (16.79)	12.40 (54.30)	8.20 (32.70)	10.98 (45.75)
Female	0.23	0.14	0.49	0.26	0.22	0.24
Married	0.61	0.78	0.23	0.49	0.53	0.48
Single	0.15	0.06	0.42	0.11	0.20	0.17
Below bachelors	0.54	0.52	0.53	0.58	0.57	0.57
Bachelors & above	0.27	0.34	0.11	0.19	0.26	0.22
Job Cat1	0.58	0.61	0.54	0.61	0.53	0.54
Job Cat2	0.11	0.12	0.07	0.10	0.12	0.12
Job Cat3	0.06	0.06	0.03	0.06	0.08	0.07
Health insurance	0.92	0.97	0.83	0.90	0.92	0.87
Race – Black	0.30	0.20	0.58	0.32	0.31	0.35
Race – Others	0.04	0.04	0.06	0.05	0.04	0.05

Head unemployed	0.06	0.03	0.12	0.07	0.06	0.08
Head NLF	0.23	0.24	0.28	0.29	0.11	0.18
West	0.19	0.19	0.18	0.20	0.19	0.20
South	0.41	0.38	0.44	0.45	0.45	0.48
Northeast	0.14	0.15	0.14	0.10	0.16	0.10
<hr/>						
Years owned						
0	18.43	—	100	—	—	—
1	4.28	—	—	18.10	14.89	17.34
2	4.57	—	—	18.40	14.10	21.39
3	4.11	—	—	19.94	14.63	13.87
4	5.50	—	—	20.55	23.67	19.94
5	7.16	—	—	23.01	32.71	27.46
6	55.96	100	—	—	—	—
<hr/>						
Observations	4092	2290	754	326	376	346
<hr/>						

Note: The first panel presents the mean and standard deviations (in parenthesis) of the continuous variables and proportions of the categorical variables in the full and subsamples. The second panel displays the column percentages for the number of years the home is owned and the third panel (i.e., last row) presents the number of observations in the full and the subsamples. NLF: Not in Labor Force.

In the sample, about 56% of individuals own a home across the entire sample period, 18% never own, and the remaining transition at least once. The age of the head of the household is that in the year 2003. Job industry is classified into four categories. *Job Cat1* is an indicator for jobs in construction, manufacturing, agriculture, and wholesale. *Job Cat2* is an indicator for jobs in business, finance, and real estate. *Job Cat3* is an indicator for jobs in the military and public services. The omitted category (*Job Cat4*) consists of jobs in professional and technical services, entertainment and arts services, health care, and other. Education is broken up into categories: less than high school (omitted), high school degree or some college (*below bachelors*), and college or advanced degree (*bachelors & above*). Race is broken up into white/Asian (omitted), *Black*, and *other*. Marital status is discretized into *married*, *single*, and *divorced/widowed* (omitted). Region is discretized to *west*, *south*, *northeast*, and *midwest* (omitted). We have two income measures, including *income-to-needs ratio* and *net wealth*.⁴ We employ an inverse hyperbolic sine (IHS) transformation for net wealth because it adjusts for skewness and retains negative and zero values, which is a common feature of data on net wealth (Friedline, Masa, & Chowa, 2015).

Table 6 demonstrates some drastic differences across the subgroups. As expected, the “owned 6 years” group is older and wealthier than the others. Families that transition tend to have more children, and a higher proportion of females and singles are in the “owned 0 years” group. These differences in the raw data motivate our question of interest – with so much state dependence in home ownership and heterogeneity among individuals and income, what are the determinants of home ownership through an economic downturn? The results should provide insights into discrepancies across subgroups of the population and should better inform policy aiming to assist home owners during downturns. Standard methods for investigating a binary panel dataset of this sort do not capture the extensive heterogeneity problem, nor do they offer quantile analyses, which highlights the usefulness of our approach.

The results for the home ownership application are presented in Table 7. Posterior means, standard deviations, and inefficiency factors calculated using the batch-means method are presented for the 25th, 50th, and 75th quantiles, as well as for the binary longitudinal probit model (PBLD). The results are based on 12,000 MCMC draws with a burn-in of 3,000 draws. The priors on the parameters are: $\beta \sim N(0_k, 10I_k)$ and $\varphi^2 \sim IG(10/2, 9/2)$. As in the female labor force application, the inefficiency factors are low, implying a nice mixing of the Markov chain.

Many of the results agree with the existing literature. Income, education, and being married all have a positive effect on home ownership (Hilber & Turner, 2014; Turner & Smith, 2009). While these align with intuition, new insights are offered across the quantiles for many of the variables. Education, for instance, is not statistically different from zero at the lower quantile. If one has a low utility for home ownership, education will not impact that decision. Additionally, age of the head has a positive impact on home ownership at the lower and median quantiles. However, for those who have a high *utility* for home ownership, age

Table 7. Results from the Home Ownership Study.

	QBLD									PBLD		
	25th Quantile			50th Quantile			75th Quantile					
	Mean	Std	IF	Mean	Std	IF	Mean	Std	IF	Mean	Std	IF
Intercept	-15.25	0.85	3.47	-9.26	0.80	4.36	-4.17	0.88	3.47	-4.15	0.31	2.27
Log age of head	1.63	0.21	3.40	0.97	0.20	3.76	0.09	0.23	3.47	0.54	0.08	2.16
No. of children	0.14	0.05	3.46	0.18	0.05	4.32	0.22	0.05	3.65	0.08	0.02	2.13
Inc-Needs ratio	0.48	0.03	8.42	0.45	0.03	8.10	0.55	0.04	6.68	0.23	0.01	6.07
IHS net wealth	0.25	0.03	4.05	0.32	0.03	4.56	0.41	0.04	5.11	0.10	0.01	2.60
Female	0.95	0.14	3.21	0.82	0.14	3.84	0.59	0.18	3.79	0.25	0.05	1.90
Married	2.28	0.14	3.42	2.18	0.15	4.05	1.70	0.17	4.35	0.74	0.05	2.05
Single	0.32	0.15	3.37	0.17	0.15	4.13	-0.27	0.17	3.49	0.01	0.05	1.97
Below bachelors	0.17	0.12	3.71	0.28	0.11	3.80	0.35	0.14	3.89	0.11	0.04	2.02
Bachelors & above	0.28	0.18	3.81	0.37	0.16	4.11	0.51	0.20	4.52	0.13	0.06	2.28
Job Cat1	0.31	0.13	4.23	0.39	0.12	4.34	0.55	0.13	4.07	0.16	0.04	2.20
Job Cat2	0.06	0.20	3.92	0.21	0.19	4.52	0.35	0.21	3.86	0.08	0.07	2.29
Job Cat3	0.03	0.24	3.54	0.08	0.23	3.82	0.11	0.26	3.74	0.01	0.08	2.04
Health insurance	0.46	0.16	3.78	0.46	0.15	3.90	0.23	0.19	4.49	0.09	0.05	2.05
Race - Black	-0.40	0.12	3.80	-0.54	0.12	3.55	-0.52	0.14	3.81	-0.18	0.04	1.96
Race - Others	-0.15	0.22	3.43	-0.51	0.21	3.80	-0.85	0.25	4.20	-0.19	0.07	1.90
Head unemployed	-0.91	0.18	3.83	-0.89	0.19	4.27	-0.76	0.23	5.13	-0.23	0.06	2.14
Head NLF	-0.40	0.14	4.03	-0.28	0.14	4.39	-0.06	0.16	4.42	-0.10	0.05	2.32
West	-0.46	0.15	3.38	-0.49	0.15	3.78	-0.52	0.18	3.95	-0.16	0.06	1.98
South	0.15	0.13	3.29	0.22	0.13	3.88	0.26	0.14	3.97	0.10	0.05	2.10
Northeast	-0.28	0.18	3.44	-0.43	0.17	3.78	-0.59	0.20	3.84	-0.17	0.06	2.15
Post-Recession (PR)	-1.32	0.23	6.22	-0.53	0.13	4.28	-0.44	0.11	2.91	-0.09	0.04	2.05
Lag-home own	7.46	0.17	5.93	5.90	0.12	4.25	9.55	0.20	7.04	2.19	0.04	2.15
PR*(lag-home own)	1.47	0.25	5.99	0.72	0.18	4.94	0.76	0.29	8.53	0.11	0.06	2.60
φ^2	0.13	0.01	9.88	0.11	0.01	8.67	0.16	0.02	8.94	0.04	0.01	8.71
Log-likelihood	-5077.07			-5030.12			-5085.64			-4446.37		
AIC	10204.14			10110.24			10221.27			8942.73		
BIC	10421.69			10327.79			10438.83			9160.29		

Note: Mean is posterior mean, Std is posterior standard deviation, and IF is inefficiency factor. NLF: Not in Labor Force.

of the head is not statistically different from zero. Number of children, however, has a positive impact across the quantiles. Family growth seems to play a role in owning a home.

The coefficient for female is positive which implies that females relative to males are more in favor of home ownership. Given that housing was previously

thought of as a safe investment, this finding aligns with [Croson and Gneezy \(2009\)](#), who investigate gender differences in preferences and find that women are more risk averse than men. Furthermore, relative to divorced/widowed individuals, being single has a positive effect only at the lower quantile. Interestingly, health insurance has a positive effect at the lower and middle quantiles and is not statistically different from zero at the higher willingness. Thus, if one has a high utility for home ownership, potential costs related to health do not play into the decision to invest in a home. While *race – Black* is negative across the quantiles, which is consistent with findings in [Charles and Hurst \(2002\)](#), *race – other* is meaningful and negative only at the middle and upper quantiles. Thus, policy interested in race disparities in home ownership should focus on high willingness individuals, because low willingness *race – other* individuals are not statistically different from whites.

The coefficient for post-Recession (2009–2013) is negative across all of the quantiles. This finding is expected given the major collapse in housing markets. The state dependence variable (*lag-home own*) is very large and positive for all of the quantiles. Even with a shock to housing markets and heterogeneity in the intercept and income controlled for, state dependence is a key element of home ownership. Interestingly, the interaction term between the state dependence variable and the post-recession indicator has a credibility interval that includes zero for the PBLD model, but is positive across the quantiles. This finding is intriguing because the positive state dependence effect offsets the negative effect from the recession. Perhaps individuals who did not own a home prior to the recession had trouble transitioning to ownership as a result of the tightened lending and credit channels. This reasoning falls in line with the work of [Hilber and Turner \(2014\)](#) in that mortgage policies can affect subgroups of home owners, but not in aggregate. The aggregate finding in PBLD shows the result is not statistically different from zero, but we find new results at the quantiles.

Covariate effect calculations, which follow from the discussion in Section 3.3, are computed for several variables in both of the models, QBLD and PBLD. The results are displayed in [Table 8](#), and show that being a female increases the probability of home ownership by 2.9 to 1.6 percentage points, for the 25th and 75th quantiles, respectively. The size of the effect is roughly halved at the 75th quantile. This is useful for understanding the differences in preferences between males and females, in particular, that at a higher willingness, they are more similar than at a lower willingness. Similar differing effects are found for the variable married, where being married increases the probability of home ownership by 8.7 percentage points at the 25th quantile and 5.4 percentage points at the 75th quantile. Furthermore, health insurance increases the probability of home ownership by 1.5 percentage points at a low willingness and 0.06 percentage points at the high willingness (although the basic result at the 75th quantile was not different from zero).

The aforementioned results find smaller effects at the higher willingness; however, this is not the case for education and wealth. Wealth and education have a greater impact for those with a high utility. Increasing net wealth by \$50,000 increases the probability of home ownership by 2.0 percentage points, and

Table 8. Covariate Effects in the Home Ownership Study.

	QBLD			PBLD
	25th	50th	75th	
Log age of head	0.0111	0.0076	0.0006	0.0128
IHS net wealth	0.0120	0.0179	0.0203	0.0177
Female	0.0298	0.0289	0.0168	0.0264
Married	0.0879	0.0890	0.0548	0.0916
Bachelors & above	0.0089	0.0132	0.0153	0.0144
Health insurance	0.0159	0.0165	0.0063	0.0098
Race – Black	–0.0133	–0.0193	–0.0149	–0.0194
Head unemployed	–0.0337	–0.0326	–0.0203	–0.0250

Note: Age is increased by 10 years and the untransformed net wealth is increased by \$50,000. The rest of the variable are indicators.

achieving a bachelors degree or more increases the probability by 1.5 percentage points. Understanding how these effects differ across the quantiles is important from a policy standpoint. For instance, if policymakers are looking to push more people into home ownership, they can consider the various types of people (high utility–low utility), and focus policy on the variables that have a greater impact on the subgroups. Additionally, when downturns occur, there are difficulties transitioning into or out of housing markets, which is clear from the results of the interaction term. These results, along with those of the demographic variables, shed light on findings that are unavailable or different than those produced from modeling the mean (PBLD).

5. CONCLUSION

This chapter presents quantile regression methods for binary longitudinal data that accommodate various forms of heterogeneity and designs an estimation algorithm to fit the model. The framework developed in this chapter contributes to literatures on quantile regression for discrete data, panel data models for quantile regression, and discrete panel data models. A simulation study is performed, which demonstrates the computational efficiency of the estimation algorithm and blocking approach.

The model is first applied to examine female labor force participation. Although this is a heavily studied topic, the panel quantile approach offers a new perspective to understand the impact of the covariates, while controlling for heterogeneity and state dependence. The results show that particular attention needs to be paid to women with newborns and children aged three to five as the impacts of these variables on female labor force participation are large and dispersed across the quantiles. The model is also applied to investigate the determinants of home ownership before, during, and after the Great Recession. The state dependence effect in home ownership is strong (even when controlling for

multivariate heterogeneity); however, after the Recession, the effect differs nontrivially from mean regression. Other results, including race, number of children, gender, health insurance, and location, also offer unique findings across the quantiles, which are unavailable in other modeling settings. The approach provided in this chapter leads to a richer view of how the covariates influence the outcome variables, which better informs policy on female labor force participation and home ownership.

NOTES

1. The derivation of the conditional posterior densities are presented in Appendix 2.
2. In the latter scenario, the model $z_i \sim N(X_i\beta + \theta w_i, \Omega_i)$ can be written as $z_i = X_i\beta + \theta w_i + L_i\eta_i$, where L_i is a lower triangular Cholesky factor of Ω_i such that $L_i L_i' = \Omega_i$. To be general, let the lower and upper truncation vectors for z_i be $a_i = (a_{i1}, \dots, a_{iT_i})$ and $b_i = (b_{i1}, \dots, b_{iT_i})$, respectively. Then the random variable η_{it} is sampled from $TN(0, 1, (a_{it} - x_{it}'\beta - \theta w_{it} - \sum_{j=1}^{t-1} l_{ij}\eta_{ij})/l_{it}, (b_{it} - x_{it}'\beta - \theta w_{it} - \sum_{j=1}^{t-1} l_{ij}\eta_{ij})/l_{it})$, where l_{ij} are the elements of L_i . This is a recursively characterized truncation region, since the range of η_{it} depends on the draw of η_{ij} for $j = 1, \dots, t-1$. The vector z_i can be obtained by substituting the recursively drawn η_i into $z_i = X_i\beta + \theta w_i + L_i\eta_i$. However, the draws so obtained are not the same as drawing z_i from a multivariate normal distribution truncated to the region $a_i < z_i < b_i$.
3. J. Chen and Ost (2005) control for state dependence in a study of housing allowance in Sweden.
4. This measure of net wealth excludes home equity and housing assets, so as to not conflate with the outcome of interest.

ACKNOWLEDGMENTS

The authors thank the anonymous referees, Ivan Jeliazkov, David Brownstone, John Geweke, K. L. Krishna, Antonio Galvao, Michael Guggisberg, and Editor Justin Tobias for their helpful comments. Discussions and suggestions from the participants at the Winter School, Delhi School of Economics (2017), and Advances in Econometrics Conference (2018) are appreciated. A special thanks to Dale Poirier for sharing the Reverend's insights and teaching us the controversy.

REFERENCES

- Albert, J., & Chib, S. (1993). Bayesian analysis of binary and polychotomous response data. *Journal of the American Statistical Association*, 88(422), 669–679.
- Albert, J. H., & Chib, S. (1996). Bayesian modeling of binary repeated measures data with application to crossover trials. In D. A. Berry & D. K. Stangl (Eds.), *Bayesian biostatistics* (pp. 577–599). New York, NY: Marcel Dekker.
- Alhamzawi, R., & Ali, H. T. M. (2018). Bayesian quantile regression for ordinal longitudinal data. *Journal of Applied Statistics*, 45(5), 815–828. doi:10.1080/02664763.2017.1315059
- Arellano, M., & Carrasco, R. (2003). Binary choice panel data models with predetermined variables. *Journal of Econometrics*, 115(1), 125–157.
- Barrodale, I., & Roberts, F. D. K. (1973). Improved algorithm for discrete l_1 linear approximation. *SIAM Journal of Numerical Analysis*, 10(5), 839–848.
- Bartolucci, F., & Farcomeni, A. (2012). A multivariate extension of the dynamic logit model for longitudinal data based on a latent Markov heterogeneity structure. *Journal of the American Statistical Association*, 104(486), 816–831.

- Bartolucci, F., & Nigro, V. (2010). A dynamic model for binary panel data with unobserved heterogeneity admitting a \sqrt{n} -consistent conditional estimator. *Econometrica*, 78(2), 719–733.
- Benoit, D. F., & Poel, D. V. D. (2012). Binary quantile regression: A Bayesian approach based on the asymmetric Laplace distribution. *Journal of Applied Econometrics*, 27(7), 1174–1188.
- Börsch-Supan, A., & Hajivassiliou, V. A. (1993). Smooth unbiased multivariate probability simulators for maximum likelihood estimation of limited dependent variable models. *Journal of Econometrics*, 58(3), 347–368.
- Breslaw, J. A. (1994). Random sampling from a truncated multivariate normal distribution. *Applied Mathematics Letters*, 7(1), 1–6.
- Burda, M., & Harding, M. (2013). Panel probit with flexible correlated effects: Quantifying technology spillovers in the presence of latent heterogeneity. *Journal of Applied Econometrics*, 28(6), 956–981.
- Carliner, G. (1974). Determinants of home ownership. *Land Economics*, 50(2), 109–119.
- Carro, J. M. (2007). Estimating dynamic panel data discrete choice models with fixed effects. *Journal of Econometrics*, 140(2), 503–528.
- Charles, K. K., & Hurst, E. (2002). The transition to home ownership and the black-white wealth gap. *The Review of Economics and Statistics*, 84(2), 281–297.
- Chen, C. (2007). A finite smoothing algorithm for quantile regression. *Journal of Computational and Graphical Statistics*, 16(1), 136–164.
- Chen, J., & Ost, C. (2005). Housing allowance and the recipient's homeownership: Evidence from a panel data study in Sweden. *Housing Studies*, 20(4), 605–625.
- Chib, S., & Carlin, B. P. (1999). On MCMC sampling in hierarchical longitudinal models. *Statistics and Computing*, 9(1), 17–26.
- Chib, S., & Greenberg, E. (1998). Analysis of multivariate probit models. *Biometrika*, 85(2), 347–361.
- Chib, S., & Jeliazkov, I. (2006). Inference in semiparametric dynamic models for binary longitudinal data. *Journal of the American Statistical Association*, 101(474), 685–700.
- Crosen, R., & Gneezy, U. (2009). Gender differences in preferences. *Journal of Economic Literature*, 47(2), 448–474.
- Dagpunar, J. (1988). *Principles of random variate generation*. Oxford: Clarendon Press.
- Dagpunar, J. (1989). An easily implemented generalized inverse Gaussian generator. *Communications in Statistics – Simulation and Computation*, 18(2), 703–710.
- Dantzig, G. B. (1963). *Linear programming and extensions*. Princeton, NJ: Princeton University Press.
- Dantzig, G. B., & Thapa, M. N. (1997). *Linear programming 1: Introduction*. New York, NY: Springer.
- Dantzig, G. B., & Thapa, M. N. (2003). *Linear programming 2: Theory and extensions*. New York, NY: Springer.
- Delattre, M., Lavielle, M., & Poursat, M.-A. (2014). A note on BIC in mixed-effects models. *Electronic Journal of Statistics*, 8(1), 456–475.
- Devroye, L. (2014). Random variate generation for the generalized inverse Gaussian distribution. *Statistics and Computing*, 24(2), 239–246.
- Drange, N., & Rege, M. (2013). Trapped at home: The effect of mothers' temporary labor market exists on their subsequent career. *Labour Economics*, 24, 125–136.
- Eckstein, Z., & Lifshitz, O. (2011). Dynamic female labor supply. *Econometrica*, 79(6), 1675–1726.
- Fairlie, R. W. (2013). Entrepreneurship, economic conditions, and the Great Recession. *Journal of Economics and Management Strategy*, 22(2), 207–231.
- Ferreira, F., Gyourko, J., & Tracy, J. (2010). Housing busts and household mobility. *Journal of Urban Economics*, 68(1), 34–45.
- Friedline, T., Masa, R. D., & Chowa, G. A. (2015). Transforming wealth: Using the inverse hyperbolic sine (ihs) and splines to predict youth's math achievement. *Social Science Research*, 49, 264–287.
- Galvao, A. F. (2010). Quantile regression for dynamic panel data with fixed effects. *Journal of Econometrics*, 164(1), 142–157.
- Galvao, A. F., & Kato, K. (2016). Smoothed quantile regression for panel data. *Journal of Econometrics*, 193(1), 92–112.

- Geraci, M., & Bottai, M. (2007). Quantile regression for longitudinal data using the asymmetric Laplace distribution. *Biostatistics*, 8(1), 140–154.
- Geweke, J. (1991). Efficient simulation from the multivariate normal and student-*t* distributions subject to linear constraints. In E. Keramidas (Ed.), *Computing science and statistics: Proceedings of the twenty-third symposium on the interface* (pp. 571–578). Fairfax: Interface Foundation of North America, Inc.
- Greenberg, E. (2012). *Introduction to Bayesian econometrics*. New York, NY: Cambridge University Press.
- Greven, S., & Kneib, T. (2010). On the behaviour of marginal and conditional AIC in linear mixed models. *Biometrika*, 97(4), 773–789.
- Hajivassiliou, V. A., & McFadden, D. L. (1998). The method of simulated scores for the estimation of LDV models. *Econometrica*, 66(4), 863–896.
- Harding, M., & Lamarche, C. (2009). A quantile regression approach for estimating panel data models using instrumental variables. *Economics Letters*, 104(3), 133–135.
- Harding, M., & Lamarche, C. (2012). Quantile regression estimation of panel duration models with censored data. *Advances in Econometrics*, 29, 237–267.
- Harding, M., & Lamarche, C. (2017). Penalized quantile regression for semiparametric models with correlated individual effects. *Journal of Applied Econometrics*, 32(2), 342–358.
- Heckman, J. J. (1981a). Heterogeneity and state dependence. In S. Rosen (Ed.), *Studies in labor markets* (pp. 91–140). Chicago, IL: University of Chicago Press.
- Heckman, J. J. (1981b). Statistical models for discrete panel data. In C. F. Manski & D. L. McFadden (Eds.), *Structural analysis of discrete data and econometric applications* (pp. 114–178). Cambridge: MIT Press.
- Heckman, J. J., & MaCurdy, T. E. (1980). A life cycle model of labor supply. *Review of Economic Studies*, 47(1), 47–74.
- Heckman, J. J., & MaCurdy, T. E. (1982). Corrigendum on a life cycle model of female labor supply. *Review of Economic Studies*, 49(4), 659–660.
- Hendricks, W., Koenker, R., & Poirier, D. (1979). Stochastic parameter models for panel data: An application to the connecticut peak load pricing experiment. *International Economic Review*, 20(3), 707–724.
- Hilber, C. A., & Turner, T. M. (2014). The mortgage interest deduction and its impact on home-ownership decisions. *The Review of Economics and Statistics*, 96(4), 618–637.
- Hsiao, C. (2007). Panel data analysis – advantages and challenges. *Test*, 16(1), 1–22.
- Hsiao, C. (2014). *Analysis of panel data (econometric society monograph)*. New York, NY: Cambridge University Press.
- Hyslop, D. R. (1999). State dependence, serial correlation and heterogeneity in intertemporal labor force participation of married women. *Econometrica*, 67(6), 1255–1294.
- Jeliaskov, I., Graves, J., & Kutzbach, M. (2008). Fitting and comparison of models for multivariate ordinal outcomes. *Advances in Econometrics: Bayesian Econometrics*, 23, 115–156.
- Jeliaskov, I., & Lee, E. (2010). MCMC perspectives on simulated likelihood estimation. *Advances in Econometrics*, 26, 3–39.
- Jeliaskov, I., & Rahman, M. A. (2012). Binary and ordinal data analysis in economics: Modeling and estimation. In X.-S. Yang (Ed.), *Mathematical modeling with multidisciplinary applications* (pp. 123–150). Hoboken, NJ: John Wiley & Sons Inc.
- Jeliaskov, I., & Vossmeier, A. (2018). The impact of estimation uncertainty on covariate effects in nonlinear models. *Statistical Papers*, 59(3), 1031–1042.
- Ji, Y., Lin, N., & Zhang, B. (2012). Model selection in binary and Tobit quantile regression using the Gibbs sampler. *Computational Statistics and Data Analysis*, 56(4), 827–839.
- Karmarkar, N. (1984). A new polynomial time algorithm for linear programming. *Combinatorica*, 4(4), 373–395.
- Keane, M. P. (1994). A computationally practical simulation estimator. *Econometrica*, 62(1), 95–116.
- Koenker, R. (2004). Quantile regression for longitudinal data. *Journal of Multivariate Analysis*, 91(1), 74–89.
- Koenker, R., & d’Orey, V. (1987). Computing regression quantiles. *Journal of the Royal Statistical Society – Series C*, 36(3), 383–393.

- Koenker, R., & Machado, J. A. F. (1999). Goodness of fit and related inference processes for quantile regression. *Journal of the American Statistical Association*, 94(448), 1296–1310.
- Koop, G., & Poirier, D. (1993). Bayesian analysis of logit models using natural conjugate priors. *Journal of Econometrics*, 56(3), 323–340.
- Koop, G., Poirier, D., & Tobias, J. (2007). *Bayesian econometric methods*. New York, NY: Cambridge University Press.
- Kordas, G. (2006). Smoothed binary regression quantiles. *Journal of Applied Econometrics*, 21(3), 387–407.
- Kotz, S., Kozubowski, T. J., & Podgorski, K. (2001). *The Laplace distribution and generalizations: A revisit with applications to communications, economics, engineering and finance*. Boston, MA: Birkhäuser.
- Kozumi, H., & Kobayashi, G. (2011). Gibbs sampling methods for Bayesian quantile regression. *Journal of Statistical Computation and Simulation*, 81(11), 1565–1578.
- Lamarche, C. (2010). Robust penalized quantile regression estimation for panel data. *Journal of Econometrics*, 157, 396–408.
- Liu, J. S. (1994). The collapsed Gibbs sampler in Bayesian computations with applications to a gene regulation problem. *Journal of the American Statistical Association*, 89(427), 958–966.
- Liu, Y., & Bottai, M. (2009). Mixed-effects models for conditional quantiles with longitudinal data. *The International Journal of Biostatistics*, 5(1), 1–22.
- Luo, Y., Lian, H., & Tian, M. (2012). Bayesian quantile regression for longitudinal data models. *Journal of Statistical Computation and Simulation*, 82(11), 1635–1649.
- Machado, J., & Silva, J. S. (2005). Quantiles for counts. *Journal of the American Statistical Association*, 100(472), 1226–1237.
- Madsen, K., & Nielsen, H. B. (1993). A finite smoothing algorithm for linear l_1 estimation. *SIAM Journal of Optimization*, 3(2), 223–235.
- Mehrotra, S. (1992). On the implementation of primal-dual interior point methods. *SIAM Journal of Optimization*, 2(4), 575–601.
- Mroz, T. M. (1987). The sensitivity of an empirical model of married women's hour of work to economic and statistical assumptions. *Econometrica*, 55(4), 765–799.
- Poirier, D. (1977). The determinants of home buying. In H. W. Watts & A. Rees (Eds.), *The New Jersey income-maintenance experiment, vol. iii: Expenditures, health, and social behaviour and the quality of the evidence* (pp. 73–91). New York, NY: Academic Press.
- Poirier, D. (2006). The growth of Bayesian methods in statistics and economics since 1970. *Bayesian Analysis*, 1(4), 969–980.
- Poirier, D., & Ruud, P. (1988). Probit with dependent observations. *Review of Economic Studies*, 55(4), 593–614.
- Portnoy, S. (2003). Censored regression quantiles. *Journal of the American Statistical Association*, 98(464), 1001–1012.
- Portnoy, S., & Koenker, R. (1997). The Gaussian hare and the Laplacian tortoise: Computability of squared-error versus absolute-error estimators. *Statistical Science*, 12(4), 279–300.
- Rahman, M. A. (2013). Quantile regression using metaheuristic algorithms. *International Journal of Computational Economics and Econometrics*, 3(3/4), 205–233.
- Rahman, M. A. (2016). Bayesian quantile regression for ordinal models. *Bayesian Analysis*, 11(1), 1–24. doi:10.1214/15-BA939
- Turner, T., & Luea, H. (2009). Homeownership, wealth accumulation, and income status. *Journal of Housing Economics*, 18, 104–114.
- Turner, T., & Smith, M. (2009). Exits from homeownership: The effects of race, ethnicity, and income. *Journal of Regional Science*, 49(1), 1–32.
- Yu, K., & Moyeed, R. A. (2001). Bayesian quantile regression. *Statistics and Probability Letters*, 54(4), 437–447.
- Yu, K., & Zhang, J. (2005). A three parameter asymmetric Laplace distribution and its extensions. *Communications in Statistics – Theory and Methods*, 34(9–10), 1867–1879.
- Yuan, Y., & Yin, G. (2010). Bayesian quantile regression for longitudinal studies with nonignorable missing data. *Biometrics*, 66(1), 105–114.

AUTHOR BIOGRAPHY

Mohammad Arshad Rahman is an Assistant Professor in the Department of Economic Sciences at the Indian Institute of Technology Kanpur, India. He obtained his PhD in Economics from the University of California, Irvine, and holds three master's degrees. His research interests include Bayesian econometrics, quantile regression, discrete choice models, Markov Chain Monte Carlo methods, computation and data analysis, and applied econometrics.

Angela Vossmeier is an Assistant Professor of Economics in the Robert Day School of Economics and Finance at Claremont McKenna College. She received her PhD in Economics in 2015 from the University of California, Irvine. Her research interests include econometrics, Bayesian inference, and financial intermediation.

APPENDIX 1 : NON-BLOCKED SAMPLING IN QBLD MODEL

The algorithm below presents the sampler for non-blocked sampling in the QBLD model.

Algorithm 2. (Non-blocked sampling)

- Let $\Psi_i = D_{\tau\sqrt{w_i}}^2$. Sample $\beta|\alpha, \varphi^2, z, w \sim N(\tilde{\beta}, \tilde{B})$, where,

$$\tilde{B}^{-1} = \left(\sum_{i=1}^n X_i' \Psi_i^{-1} X_i + B_0^{-1} \right) \text{ and}$$

$$\tilde{\beta} = \tilde{B} \left(\sum_{i=1}^n X_i' \Psi_i^{-1} (z_i - S_i \alpha_i - \theta w_i) + B_0^{-1} \beta_0 \right).$$

- Sample $\alpha_i|\beta, \varphi^2, z, w \sim N(\tilde{\alpha}, \tilde{A})$ for $i = 1, \dots, n$, where,

$$\tilde{A}^{-1} = \left(S_i' D_{\tau\sqrt{w_i}}^{-2} S_i + \frac{1}{\varphi^2} I_l \right) \text{ and } \tilde{\alpha} = \tilde{A} \left(S_i' D_{\tau\sqrt{w_i}}^{-2} (z_i - X_i \beta - \theta w_i) \right).$$

- Sample $w_{it}|\beta, \alpha_i, z_{it} \sim GIG(0.5, \tilde{\lambda}_{it}, \tilde{\eta})$ for $i = 1, \dots, n$ and $t = 1, \dots, T_i$, where,

$$\tilde{\lambda}_{it} = \left(\frac{z_{it} - x'_{it}\beta - s'_{it}\alpha_i}{\tau} \right)^2 \text{ and } \tilde{\eta} = \left(\frac{\theta^2}{\tau^2} + 2 \right).$$

- Sample $\varphi^2|\alpha \sim IG(\tilde{c}_1/2, \tilde{d}_1/2)$, where $\tilde{c}_1 = (nl + c_1)$ and $\tilde{d}_1 = (\sum_i \alpha_i' \alpha_i + d_1)$.
- Sample the latent variable $z|y, \beta, \alpha, w$ for all values of $i = 1, \dots, n$ and $t = 1, \dots, T_i$ from an univariate truncated normal (TN) distribution as follows,

$$z_{it}|y, \beta, w \sim \begin{cases} TN_{(-\infty, 0]}(x'_{it}\beta + s'_{it}\alpha_i + \theta w_{it}, \tau^2 w_{it}) & \text{if } y_{it} = 0, \\ TN_{(0, \infty)}(x'_{it}\beta + s'_{it}\alpha_i + \theta w_{it}, \tau^2 w_{it}) & \text{if } y_{it} = 1. \end{cases}$$

APPENDIX 2 : THE CONDITIONAL DENSITIES FOR BLOCKED SAMPLING IN QBLD MODEL

This appendix presents a derivation of the conditional posterior densities for blocked sampling in the QBLD model. Specifically, the parameters β and latent variable z_i are sampled marginally of the random effects parameter α_i from an updated multivariate normal and a truncated multivariate normal distribution, respectively. The parameter α_i is sampled conditional on (β, z_i) from an updated multivariate normal distribution. The latent weights w are sampled element-wise from a GIG distribution and the variance φ^2 is sampled from an updated IG distribution.

- The mean and variance of the QBLD model, $z_i = X_i\beta + S_i\alpha_i + \theta w_i + D_{\tau_i\sqrt{w_i}}u_i$ for $i = 1, \dots, n$, (marginally of α_i) can be shown to have the following expressions,

$$\begin{aligned} E(z_i) &= X_i\beta + \theta w_i, \\ V(z_i) &= \varphi^2 S_i S_i' + D_{\tau_i\sqrt{w_i}}^2 = \Omega_i. \end{aligned}$$

First, we derive the conditional posterior of β and z_i , marginally of α_i , but conditional on other variables in the model.

- Starting with β , the conditional posterior density $\pi(\beta|z, w, \varphi^2)$ can be derived as,

$$\begin{aligned} \pi(\beta|z, w, \varphi^2) &\propto \left\{ \prod_{i=1}^n f(z_i|\beta, w_i, \varphi^2) \right\} \pi(\beta) \\ &\propto \exp \left[-\frac{1}{2} \left\{ \sum_{i=1}^n (z_i - X_i\beta - \theta w_i)' \Omega_i^{-1} (z_i - X_i\beta - \theta w_i) \right. \right. \\ &\quad \left. \left. + (\beta - \beta_0)' B_0^{-1} (\beta - \beta_0) \right\} \right] \\ &\propto \exp \left[-\frac{1}{2} \left\{ \beta' \left(\sum_{i=1}^n X_i' \Omega_i^{-1} X_i + B_0^{-1} \right) \beta \right. \right. \\ &\quad \left. \left. - \beta' \left(\sum_{i=1}^n X_i' \Omega_i^{-1} (z_i - \theta w_i) + B_0^{-1} \beta_0 \right) \right. \right. \\ &\quad \left. \left. - \left(\sum_{i=1}^n (z_i - \theta w_i)' \Omega_i^{-1} X_i + \beta_0' B_0^{-1} \right) \beta \right\} \right] \\ &\propto \exp \left[-\frac{1}{2} \left\{ \beta' \tilde{B}^{-1} \beta - \beta' \tilde{B}^{-1} \tilde{\beta} - \tilde{\beta}' \tilde{B}^{-1} \beta \right\} \right], \end{aligned}$$

where the third line only keeps terms involving β and the fourth line introduces the terms $\tilde{\beta}$ and \tilde{B} , which are defined as,

$$\tilde{B}^{-1} = \left(\sum_{i=1}^n X_i' \Omega_i^{-1} X_i + B_0^{-1} \right) \quad \text{and} \quad \tilde{\beta} = \tilde{B} (X_i' \Omega_i^{-1} (z_i - \theta w_i) + B_0^{-1} \beta_0).$$

Adding and subtracting $\tilde{\beta}' \tilde{B}^{-1} \tilde{\beta}$ and absorbing the term $\exp \left[-\frac{1}{2} \left\{ -\tilde{\beta}' \tilde{B}^{-1} \tilde{\beta} \right\} \right]$ into the proportionality constant, the square can be completed as follows,

$$\pi(\beta|z, w, \varphi^2) \propto \exp \left[-\frac{1}{2} (\beta - \tilde{\beta})' \tilde{B}^{-1} (\beta - \tilde{\beta}) \right].$$

The above expression is recognized as the kernel of a Gaussian or normal distribution and hence $\beta|z, w, \varphi^2 \sim N(\tilde{\beta}, \tilde{B})$.

- The conditional posterior density of the latent variable z marginally of α can be obtained from the joint posterior density Eq. (8) as,

$$\begin{aligned} \pi(z|\beta, w, \varphi^2, y) &\propto \prod_{i=1}^n \{ \pi(z_i|\beta, w_i, \varphi^2, y_i) \} \\ &\propto \prod_{i=1}^n \left\{ \prod_{t=1}^{T_i} [I(z_{it} > 0)I(y_{it} = 1) + I(z_{it} \leq 0)I(y_{it} = 0)] \right. \\ &\quad \left. \times \exp \left[-\frac{1}{2} (z_i - X_i \beta - \theta w_i)' \Omega_i^{-1} (z_i - X_i \beta - \theta w_i) \right] \right\}. \end{aligned}$$

The expression inside the curly braces corresponds to a truncated multivariate normal distribution, so $z_i|y_i, \beta, w_i, \varphi^2 \sim TMVNB_{B_i}(X_i \beta + \theta w_i, \Omega_i)$ for all $i = 1, \dots, n$. Here, B_i is the truncation region such that $B_i = (B_{i1} \times B_{i2} \times \dots \times B_{iT_i})$, where B_{it} is the interval $(0, \infty)$ if $y_{it} = 1$ and the interval $(-\infty, 0]$ if $y_{it} = 0$ for $t = 1, \dots, T_i$. Sampling directly from a truncated multivariate normal distribution (TMVN) is not possible, hence we resort to the method proposed in Geweke (1991), which utilizes Gibbs sampling to make draws from a TMVN.

Let z_i^j denote the values of z_i at the j -th pass of the MCMC iteration. Then sampling is done from a series of conditional posterior distribution as follows:

$$z_{it}^j | z_{i1}^j, \dots, z_{i(t-1)}^j, z_{i(t+1)}^j, \dots, z_{iT_i}^j \sim TN_{B_{it}}(\mu_{\ell-t}, \Sigma_{\ell-t}), \quad \text{for } t = 1, \dots, T_i,$$

where TN denotes a truncated normal distribution. The terms $\mu_{\ell-t}$ and $\Sigma_{\ell-t}$ are the conditional mean and variance, respectively, and are defined as,

$$\begin{aligned} \mu_{\ell-t} &= x'_{it} \beta + \theta w_{it} + \Sigma_{t,-t} \Sigma_{-t,-t}^{-1} (z_{i,-t}^j - (X_i \beta + \theta w_i)_{-t}), \\ \Sigma_{\ell-t} &= \Sigma_{t,t} - \Sigma_{t,-t} \Sigma_{-t,-t}^{-1} \Sigma_{-t,t}, \end{aligned}$$

where $z_{i,-t}^j = (z_{i1}^j, \dots, z_{i(t-1)}^j, z_{i(t+1)}^j, \dots, z_{iT_i}^j)$, $(X_i \beta + \theta w_i)_{-t}$ is a column vector with t -th element removed, $\Sigma_{t,t}$ denotes the (t, t) -th element of Ω_i , $\Sigma_{t,-t}$

denotes the t -th row of Ω_i with element in the t -th column removed, and $\Sigma_{-t,-t}$ is the Ω_i matrix with t -th row and t -th column removed.

- The conditional posterior density of the random effects parameters α_i for $i = 1, \dots, n$ is derived from the joint posterior density (8) as follows,

$$\begin{aligned} \pi(\alpha_i | z_i, \beta, w_i, \varphi^2) &\propto f(z_i | \beta, \alpha_i, w_i) \pi(\alpha_i | \varphi^2) \\ &\propto \exp \left[-\frac{1}{2} \left\{ (z_i - X_i \beta - S_i \alpha_i - \theta w_i)' D_{\tau \sqrt{w_i}}^{-2} (z_i - X_i \beta - S_i \alpha_i - \theta w_i) \right. \right. \\ &\quad \left. \left. + \frac{\alpha_i' \alpha_i}{\varphi^2} \right\} \right] \\ &\propto \exp \left[-\frac{1}{2} \left\{ \alpha_i' (S_i' D_{\tau \sqrt{w_i}}^{-2} S_i + \varphi^{-2} I_l) \alpha_i - \alpha_i' (S_i' D_{\tau \sqrt{w_i}}^{-2} (z_i - X_i \beta - \theta w_i)) \right. \right. \\ &\quad \left. \left. - ((z_i - X_i \beta - \theta w_i)' D_{\tau \sqrt{w_i}}^{-2} S_i) \alpha_i \right\} \right] \\ &\propto \exp \left[-\frac{1}{2} (\alpha_i - \tilde{a})' \tilde{A}^{-1} (\alpha_i - \tilde{a}) \right], \end{aligned}$$

where the third line omits all terms not involving α_i and the fourth line introduces the terms,

$$\tilde{A}^{-1} = \left(S_i' D_{\tau \sqrt{w_i}}^{-2} S_i + \frac{1}{\varphi^2} I_l \right) \quad \text{and} \quad \tilde{a} = \tilde{A} \left(S_i' D_{\tau \sqrt{w_i}}^{-2} (z_i - X_i \beta - \theta w_i) \right),$$

as the posterior precision and posterior mean, respectively, and completes the square. The result is a kernel of a normal distribution, hence, $\alpha_i | z_i, \beta, w_i, \varphi^2 \sim N(\tilde{a}, \tilde{A})$ for $i = 1, \dots, n$.

- The conditional posterior density of w is obtained from the joint posterior density (8) by collecting terms involving w . Each term in w is updated element-wise as follows:

$$\begin{aligned} \pi(w_{it} | z_{it}, \beta, \alpha_i) &\propto (2\pi\tau^2 w_{it})^{-1/2} \exp \left[-\frac{1}{2\tau^2 w_{it}} (z_{it} - x'_{it} \beta - s'_{it} \alpha_i - \theta w_{it})^2 - w_{it} \right] \\ &\propto w_{it}^{-1/2} \exp \left[-\frac{1}{2} \left\{ \left(\frac{z_{it} - x'_{it} \beta - s'_{it} \alpha_i}{\tau} \right)^2 w_{it}^{-1} + \left(\frac{\theta^2}{\tau^2} + 2 \right) w_{it} \right\} \right] \\ &\propto w_{it}^{-1/2} \exp \left[-\frac{1}{2} \{ \tilde{\lambda}_{it} w_{it}^{-1} + \tilde{\eta} w_{it} \} \right], \end{aligned}$$

where the second line omits all terms not involving w_{it} and the third line introduces the terms defined below,

$$\tilde{\lambda}_{it} = \left(\frac{z_{it} - x'_{it}\beta - s'_{it}\alpha_i}{\tau} \right)^2 \quad \text{and} \quad \tilde{\eta} = \left(\frac{\theta^2}{\tau^2} + 2 \right).$$

The expression in the third line is recognized as the kernel of a GIG distribution. Hence, we have $w_{it}|z_{it}, \beta, \alpha_i \sim GIG(0.5, \tilde{\lambda}_{it}, \tilde{\eta})$ for $t = 1, \dots, T_i$ and $i = 1, \dots, n$.

- The conditional posterior density of φ^2 is obtained from the joint posterior density (8) by collecting terms involving φ^2 conditional on the remaining model parameters. This is done below.

$$\begin{aligned} \pi(\varphi^2|\alpha) \sim \dots &\propto (2\pi)^{-nl/2} (\varphi^2)^{-nl/2} \exp \left[-\frac{1}{2\varphi^2} \sum_{i=1}^n \alpha'_i \alpha_i \right] (\varphi^2)^{-(c_1/2+1)} \exp \left[-\frac{d_1}{2\varphi^2} \right] \\ &\propto (\varphi^2)^{-(nl/2+c_1/2+1)} \exp \left[-\frac{1}{2\varphi^2} \left\{ \sum_{i=1}^n \alpha'_i \alpha_i + d_1 \right\} \right] \\ &\propto (\varphi^2)^{(\tilde{c}_1/2+1)} \exp \left[-\frac{1}{2\varphi^2} \tilde{d}_1 \right], \end{aligned}$$

where $\tilde{c}_1 = nl + c_1$ and $\tilde{d}_1 = (\sum_{i=1}^n \alpha'_i \alpha_i + d_1)$. The expression in the last line is recognized as the kernel of an IG distribution and consequently, we have $\varphi^2|\alpha \sim IG(\tilde{c}_1/2, \tilde{d}_1/2) \sim IG(\tilde{c}_1/2, \tilde{d}_1/2)$.